


### **Synonym**

HPAD10, PAD1, PDI, PDI1

#### **Source**

Human PADI1 Protein, His Tag(PA1-H5543) is expressed from Baculovirus-Insect cells. It contains AA Met 1 - Pro 663 (Accession # Q9ULC6). Predicted N-terminus: Met

### **Molecular Characterization**



This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 76.7 kDa. The protein migrates as 47 kDa and 65-70 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### **Endotoxin**

Less than 1.0 EU per  $\mu g$  by the LAL method.

# **Purity**

>95% as determined by SDS-PAGE.

#### **Formulation**

Supplied as 0.2  $\mu m$  filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with glycerol as protectant.

Contact us for customized product form or formulation.

## **Shipping**

This product is supplied and shipped with dry ice, please inquire the shipping cost.


### **Storage**

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

### **SDS-PAGE**



Human PADI1 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

### **Background**

This gene encodes a member of the peptidyl arginine deiminase family of enzymes, which catalyze the post-translational deimination of proteins by converting arginine residues into citrullines in the presence of calcium ions. The family members have distinct substrate specificities and tissue-specific expression patterns. The type I enzyme is involved in the late stages of epidermal differentiation, where it deiminates filaggrin and keratin K1, which maintains hydration of the stratum corneum, and hence the cutaneous barrier function. This enzyme may also play a role in hair follicle formation. This gene exists in a cluster with four other paralogous genes. [provided by RefSeq, Jul 2008]

# **Clinical and Translational Updates**

