Monoclonal Anti-RSV-post-F0 specific Antibody, Mouse IgG1 (1A2D9D8)





Source

Monoclonal Anti-RSV-post-F0 specific Antibody, Mouse IgG1 (1A2D9D8) is a Mouse monoclonal antibody produced from a hybridoma created by fusing SP2/0 myeloma and Mouse B-lymphocytes.

Clone

1A2D9D8

Species

Mouse

Isotype

Mouse IgG1 | Mouse Kappa

Conjugate

Unconjugated

Antibody Type

Hybridoma Monoclonal

Reactivity

Virus

Immunogen

Recombinant HRSV (A) Fusion glycoprotein F0 derived from human 293 cells.

Specificity

This product is a specific antibody specifically reacts with Postfusion glycoprotein F0/post-F protein (RSV).

Application

Application Recommended Usage

ELISA 0.5-500 ng/mL

Purity

>95% as determined by SDS-PAGE.

Purification

Protein A purified/ Protein G purified

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Monoclonal Anti-RSV-post-F0 specific Antibody, Mouse IgG1 (1A2D9D8)








Monoclonal Anti-RSV-post-F0 specific Antibody, Mouse IgG1 (1A2D9D8) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With <u>Star Ribbon Prestained Protein Marker</u>).

Bioactivity-ELISA



Immobilized HRSV (A) Post-fusion glycoprotein F0, His Tag (MALS verified) (Cat. No. RSF-V52H6) at 2 μ g/mL (100 μ L/well) can bind Monoclonal Anti-RSV-post-F0 specific Antibody, Mouse IgG1 (1A2D9D8) (Cat. No. RSV-Y180) with a linear range of 0.488-62.5 ng/mL. The antibody does not bind HRSV (A) Pre-fusion glycoprotein F0, His Tag (MALS verified) (Cat. No. RSF-V52H7) (QC tested).

Background

Human respiratory syncytial virus (HRSV) is the most common etiological agent of acute lower respiratory tract disease in infants and can cause repeated infections throughout life. The RSV fusion glycoprotein (RSV F) is the principal target of RSV neutralizing antibodies in human sera. The RSV F is a type I viral fusion protein synthesized as inactive, single-chain polypeptides that assemble into trimers. RSV F fuses the viral and host cell membranes by irreversible protein refolding from the labile prefusion conformation to the stable post-fusion conformation.

Clinical and Translational Updates

