

## **Synonym**

FLJ18683,T3E,TCRE,CD3E,CD3-epsilon

## **Source**

Cynomolgus CD3 epsilon, His Tag(CDE-C5226) is expressed from human 293 cells (HEK293). It contains AA Gln 22 - Asp 117 (Accession # Q95LI5-1). Predicted N-terminus: Gln 22

#### **Molecular Characterization**

CD3 epsilon(Gln 22 - Asp 117) Q95LI5-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 16.1 kDa. The protein migrates as 18-23 kDa under reducing (R) condition, and 33-40 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

#### Endotoxin

Less than 1.0 EU per µg by the LAL method.

# **Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


## **SDS-PAGE**



Cynomolgus CD3 epsilon, His Tag on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

## **Bioactivity-ELISA**


## **SEC-MALS**



The purity of Cynomolgus CD3 epsilon, His Tag (Cat. No. CDE-C5226) is more than 90% and the molecular weight of this protein is around 35-45 kDa verified by SEC-MALS.


Report






Immobilize Cynomolgus CD3 epsilon, His Tag (Cat. No. CDE-C5226) at 2  $\mu$ g/mL, add increasing concentrations of Bispecific T cell Engager (CD3 X BCMA), and then add Biotinylated Human BCMA, Fc,Avitag (Cat. No. BC7-H82F0) at 0.2  $\mu$ g/mL. Detection is performed using HRP-conjugated streptavidin. Binding occurs at a linear range of 1-5 ng/mL (QC tested).

# **Bioactivity-BLI**



Loaded Anti-Human CD3 mAb, mouse IgG1 (Clone # SP3541-2) on AMC Biosensor, can bind Cynomolgus CD3 epsilon, His Tag (Cat. No. CDE-C5226) with an affinity constant of 0.305 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Bispecific T-cell Engager (CD3 X BCMA) on AHC Biosensor via DMF Filed Human BCMA, Fc Tag (Cat. No. BC7-H525541), can bind Cynomolgus CD3 epsilon, His Tag (Cat. No. CDE-C5226) with an affinity constant of 0.113 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

# Background

CD3e molecule, epsilon is also known as CD3E, is a T-cell surface single-pass type I membrane glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3E, together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development, and defects in CD3E gene cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to type I diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.

# **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.

