

**Synonym**


IL-17C,Cytokine CX2,IL17C

**Source**

Biotinylated Mouse IL-17C, His,Avitag(ILC-M82E9) is expressed from human 293 cells (HEK293). It contains AA Asp 17 - Gln 194 (Accession # [Q8K4C5-1](#) ).

Predicted N-terminus: Asp 17

**Molecular Characterization**



This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™)

The protein has a calculated MW of 23.4 kDa. The protein migrates as 25-27 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Endotoxin**

Less than 1.0 EU per µg by the LAL method.

**Purity**

>95% as determined by SDS-PAGE.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, 0.5 M Arginine, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

**Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


**SDS-PAGE**



Biotinylated Mouse IL-17C, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


**Bioactivity-ELISA**

**SEC-MALS**



The purity of Biotinylated Mouse IL-17C, His,Avitag (Cat. No. ILC-M82E9) is more than 85% and the molecular weight of this protein is around 57-67 kDa verified by SEC-MALS.

[Report](#)



Immobilized Human IL-17 RE (155-454), Fc Tag (Cat. No. ILE-H5256) at 2 µg/mL (100 µL/well) can bind Biotinylated Mouse IL-17C, His,Avitag (Cat. No. ILC-M82E9) with a linear range of 0.2-20 ng/mL (QC tested).

### Background

Interleukin-17C(IL-17C) is a glycosylated cytokine that plays an important role in mucosal immunity and chronic inflammation. IL-17C binds to IL-17 RE with high affinity and to IL-17 RA with low affinity, binds to a heterodimer formed by IL17RA and IL17RE. Enhanced IL17C/IL17RE signaling may also lead to greater susceptibility to autoimmune diseases. Stimulates the production of antibacterial peptides and proinflammatory molecules for host defense by signaling through the NF-kappa-B and MAPK pathways. Acts synergically with IL22 in inducing the expression of antibacterial peptides, including S100A8, S100A9, REG3A and REG3G. Synergy is also observed with TNF and IL1B in inducing DEFB2 from keratinocytes.

### Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.