



## Synonym

Integrin alpha V beta 1,ITGAV&ITGB1

## Source

Human ITGAV&ITGB1 Heterodimer Protein, His Tag&Tag Free(IT1-H52E1) is expressed from human 293 cells (HEK293). It contains AA Phe 31 - Val 992 (ITGAV) & Gln 21 - Asp 728 (ITGB1) (Accession # [NP\\_002201.1](#) (ITGAV) & [NP\\_002202.2](#) (ITGB1)).

Predicted N-terminus: Phe 31 (ITGAV) & Gln 21 (ITGB1)

## Molecular Characterization

ITGAV (Phe 31 - Val 992) NP_002201.1	Acidic Tail	Poly-his
ITGB1 (Gln 21 - Asp 728) NP_002202.2	Basic Tail	

Human ITGAV&ITGB1 Heterodimer Protein, His Tag&Tag Free, produced by co-expression of ITGAV and ITGB1, has a calculated MW of 112.9 kDa (ITGAV) and 83.7 kDa (ITGB1). Subunit ITGAV is fused with an acidic tail at the C-terminus and followed by a polyhistidine tag and subunit ITGB1 contains no tag but a basic tail at the C-terminus. The non-reducing (NR) protein migrates as 125-150 kDa (ITGAV) and 95-110 kDa (ITGB1) respectively due to glycosylation.

## Endotoxin

Less than 1.0 EU per µg by the LAL method.

## Purity

>90% as determined by SDS-PAGE.

## Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE




Human ITGAV&ITGB1 Heterodimer Protein, His Tag&Tag Free on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## Bioactivity-ELISA

Discounts, Gifts,  
and more!





Immobilized Human Fibronectin at 2 µg/mL (100 µL/well) can bind Human ITGAV&ITGB1 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT1-H52E1) with a linear range of 0.002-0.063 µg/mL (QC tested).

## Background

Integrin alpha-5/beta-1 is a receptor for fibrinogen. Integrin alpha-1/beta-1, alpha-2/beta-1, alpha-6/beta-1 and alpha-7/beta-1 are receptors for laminin. Integrin alpha-4/beta-1 is a receptor for VCAM1. It recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-9/beta-1 is a receptor for VCAM1, cytotactin and osteopontin. It recognizes the sequence A-E-I-D-G-I-E-L in cytotactin. Integrin alpha-V/beta-1 is also a receptor for vitronectin. Beta-1 integrins recognize the sequence R-G-D in a wide array of ligands. Isoform 2 interferes with isoform 1 resulting in a dominant negative effect on cell adhesion and migration (in vitro). When associated with alpha-7/beta-1 integrin, regulates cell adhesion and laminin matrix deposition.

## Clinical and Translational Updates

Discounts, Gifts,  
and more!

