

### Synonym

Spike,S2 protein,Spike glycoprotein Subunit2,S glycoprotein Subunit2,Spike protein S2

### Source

SARS-CoV-2 S2 protein (A701V), His Tag (S2N-C52Hc) is expressed from human 293 cells (HEK293). It contains AA Ser 686 - Pro 1213 (Accession # [QHD43416.1](#)). The recombinant protein is expressed from human 293 cells (HEK293) with T4 fibrin trimerization motif and a polyhistidine tag at the C-terminus. Proline substitutions (F817P/ A892P/ A899P/ A942P/ K986P/ V987P) are introduced to stabilize the trimeric prefusion state of SARS-CoV-2 S protein. The A701V mutation was identified in the SARS-CoV-2 Beta variant (Pango lineage: B.1.351; other names: 20H/501Y.V2).

Predicted N-terminus: Ser 686

### Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus

The protein has a calculated MW of 59.8 kDa. The protein migrates as 70-100 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per µg by the LAL method.

### Purity

>95% as determined by SDS-PAGE.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

### SDS-PAGE




SARS-CoV-2 S2 protein (A701V), His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

### Bioactivity-ELISA



Immobilized SARS-CoV-2 S2 protein (A701V), His Tag (Cat. No. S2N-C52Hc) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike S2 protein Antibody, Human IgG4 (Cat. No. S2N-S86) with a linear range of 0.2-2 ng/mL (QC tested).



Immobilized Anti-SARS-CoV-2 Spike S2 protein Antibody, Human IgG4 (Cat. No. S2N-S86) at 1 µg/mL (100 µL/well) can bind SARS-CoV-2 S2 protein (A701V), His Tag (Cat. No. S2N-C52Hc) with a linear range of 0.8-13 ng/mL (Routinely tested).

## Background

It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.